Duilding blocks for MI immunity

Recommended vaccines for children from birth through 6 years old

RV

Rotavirus

Hib

Hib meningitis

PCV

Pneumococcal

IPV

Polio

DTaP

Diphtheria,

Tetanus, Pertussis

AGF 4

MONTHS

Timely vaccination is recommended by the Centers for Disease Control Prevention, the American Academy of Pediatrics and the American Academy of Family Physicians and are endorsed by the Michigan Department of Health and Human Services



Ask your health care provider about combination vaccines to reduce the number of pokes in a visit.

*The number of doses in the series will depend on the vaccine product that your health care provider carries.



BIRTH

HepB Hepatitis B

> RV Rotavirus

Hib Hib meningitis

PCV Pneumococcal

> **IPV** Polio

DTaP Diphtheria, Tetanus, Pertussis

AGE 2 **MONTHS** Not earlier than 6 weeks of age

HepB Hepatitis B

*RV Rotavirus

*Hib Hib meningitis

PCV Pneumococca

IPV Polio

DTaP Diphtheria, Tetanus, Pertussis

> AGF 6 **MONTHS**

Varicella Chickenpox

Hib Hib meningitis

PCV Pneumococcal

MMR Measles, Mumps. Rubella

> HepA Hepatitis A

DTaP

Diphtheria,

Tetanus, Pertussis

AGF 15

MONTHS

AGE 12 **MONTHS** On or after first birthday

Everyone age 6 months and older is recommended to aet flu vaccine each year.

Children 6 months through 8 years old may need 2 doses of vaccine, given 4 weeks apart. Ask your health care provider about your child's vaccine needs.

HepA

Hepatitis A

AGE 18

MONTHS









Diphtheria, Tetanus, Pertussis

> AGE 4 TO 6 YEARS Before Kindergarten

The Michigan Department of Health and Human Services will not exclude from participation in, deny benefits of, or discriminate against any individual or group because of race, sex, religion, age, national origin, color, height, weight, marital status, gender identification or expression, sexual orientation, partisan considerations, or a disability or genetic information that is unrelated to the person's eligibility.